# ABclonal®

## Von Hippel Lindau/VHL Rabbit mAb

Catalog No.: A23239 Recombinant 1 Publications

## **Basic Information**

## **Observed MW**

18-24kDa

#### **Calculated MW**

24kDa

#### Category

SMab Recombinant Monoclonal Antibody

## **Applications**

WB, ELISA

## **Cross-Reactivity**

Human, Mouse, Rat

#### CloneNo number

ARC59984

## **Background**

This gene encodes a component of a ubiquitination complex. The encoded protein is involved in the ubiquitination and degradation of hypoxia-inducible-factor (HIF), which is a transcription factor that plays a central role in the regulation of gene expression by oxygen. In addition to oxygen-related gene expression, this protein plays a role in many other cellular processes including cilia formation, cytokine signaling, regulation of senescence, and formation of the extracellular matrix. Variants of this gene are associated with von Hippel-Lindau syndrome, pheochromocytoma, erythrocytosis, renal cell carcinoma, and cerebellar hemangioblastoma.

## **Recommended Dilutions**

WB 1:500 - 1:1000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID Swiss Prot** 7428 P40337

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 100-200 of human Von Hippel Lindau/VHL. (NP\_000542.1).

#### **Synonyms**

RCA1; VHL1; pVHL; HRCA1; Von Hippel Lindau/VHL

## **Contact**

www.abclonal.com

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

## Validation Data



Western blot analysis of various lysates, using Von Hippel Lindau/VHL Rabbit mAb (A23239) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 45s.