A2352

Leader in Biomolecular Solutions for Life Science

www.abclonal.com

Histone H3 Rabbit pAb

Catalog No.: A2352 10 Publications

Basic Information

Observed MW
17 kDa

Calculated MW
15 kDa

Category
Polyclonal Antibody

Applications
WB,IHC-P,IP,ChIP,ELISA

Cross-Reactivity
Human,Mouse,Rat,Other (Wide Range Predicted)

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Recommended Dilutions

WB	1:2000 - 1:10000
IHC-P	1:50 - 1:200
IP	0.5μg-4μg antibody for 200μg-400μg extracts of whole cells
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.
ChIP	5μg antibody for 5μg-10μg of Chromatin

Immunogen Information

Gene ID
8290/8350

Swiss Prot
Q16695/P68431

Immunogen
Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms
H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3; Histone H3

Product Information

Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20℃


Contact

www.abclonal.com


Validation Data




Western blot analysis of various lysates using Histone H3 Rabbit pAb (A2352) at 1:10000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 60s.




Immunohistochemistry analysis of paraffin-embedded Human breast cancer using Histone H3 Rabbit pAb (A2352) at dilution of 1:20 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse liver using Histone H3 Rabbit pAb (A2352) at dilution of 1:20 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Rat ovary using Histone H3 Rabbit pAb (A2352) at dilution of 1:20 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunoprecipitation analysis of 150 µg extracts of MCF7 cells using 3 µg Histone H3 antibody (A2352). Western blot was performed from the immunoprecipitate using Histone H3 antibody (A2352) at a dilution of 1:500.

Validation Data



Chromatin immunoprecipitation analysis of extracts of HeLa cells, using Histone H3 antibody (A2352) and rabbit IgG.The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.

For research use only. Not for therapeutic or diagnostic purposes.
Please visit https://abclonal.com for a complete listing of recommended products.