A2355SP

Leader in Biomolecular Solutions for Life Science

www.abclonal.com

MonoMethyl-Histone H3-K4 Rabbit pAb

Catalog No.: A2355SP 41 Publications

Basic Information

Observed MW
17 kDa

Calculated MW
15 kDa

Category
Polyclonal Antibody

Applications
WB,IHC-P,IF/ICC,ChIP,ELISA,DB

Cross-Reactivity
Human,Mouse,Rat,Other (Wide Range Predicted)

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Recommended Dilutions

WB	1:1000 - 1:5000
IF/ICC	1:200 - 1:800
IHC-P	1:2000 - 1:8000
DB	1:1000 - 1:2000
ChIP	3 μg antibody for 5 μg - 10 μg of Chromatin
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.For high-ratio antibody dilutions (≥1:10000) , a sequential dilution method is strongly recommended to ensure measurement accuracy.

Immunogen Information

Gene ID
8290/8350

Swiss Prot
Q16695/P68431

Immunogen
Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms
H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3; MonoMethyl-Histone H3-K4

Product Information

Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20℃


Contact

www.abclonal.com


Validation Data




Western blot analysis of various lysates using MonoMethyl-Histone H3-K4 Rabbit pAb (A2355SP) at 1:5000 dilution incubated overnight at 4℃.|Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.|Lysates/proteins: 25 µg per lane.|Blocking buffer: 3% nonfat dry milk in TBST.|Detection: ECL Basic Kit (RM00020).|Negative control (NC): H3 protein.|Exposure time: 90 s




Chromatin immunoprecipitation was performed with 10 µg of cross-linked chromatin from HeLa cells, using 3 µg of MonoMethyl-Histone H3-K4 Rabbit pAb (A2355SP) and Rabbit IgG isotype control (AC042). The enrichment of immunoprecipitated DNA at different genomic loci was examined by quantitative PCR. The histogram compares the ratio of the immunoprecipitated DNA to the input at given loci




Immunohistochemistry analysis of paraffin-embedded Mouse kidney tissue using MonoMethyl-Histone H3-K4 Rabbit pAb (A2355SP) at a dilution of 1:4000 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining





Immunohistochemistry analysis of paraffin-embedded Mouse lung tissue using MonoMethyl-Histone H3-K4 Rabbit pAb (A2355SP) at a dilution of 1:4000 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining



Immunohistochemistry analysis of paraffin-embedded Rat colon tissue using MonoMethyl-Histone H3-K4 Rabbit pAb (A2355SP) at a dilution of 1:4000 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining








Validation Data

Dot-blot analysis of all sorts of peptides using MonoMethyl-Histone H3-K4 Rabbit pAb (A2355SP) at 1:1000 dilution

Confocal imaging of C6 cells using MonoMethyl-Histone H3-K4 Rabbit pAb (A2355SP, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x

Confocal imaging of HeLa cells using MonoMethyl-Histone H3-K4 Rabbit pAb (A2355SP, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x



Confocal imaging of NIH/3T3 cells using MonoMethyl-Histone H3-K4 Rabbit pAb (A2355SP, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x