ABclonal www.abclonal.com

hnRNP E2/PCBP2 Rabbit mAb

Catalog No.: A23987 Recombinant 1 Publications

Basic Information

Observed MW

35kDa/45kDa

Calculated MW

39kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

WB,IP,ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC56507

Background

The protein encoded by this gene appears to be multifunctional. Along with PCBP-1 and hnRNPK, it is one of the major cellular poly(rC)-binding proteins. The encoded protein contains three K-homologous (KH) domains which may be involved in RNA binding. Together with PCBP-1, this protein also functions as a translational coactivator of poliovirus RNA via a sequence-specific interaction with stem-loop IV of the IRES, promoting poliovirus RNA replication by binding to its 5'-terminal cloverleaf structure. It has also been implicated in translational control of the 15-lipoxygenase mRNA, human papillomavirus type 16 L2 mRNA, and hepatitis A virus RNA. The encoded protein is also suggested to play a part in formation of a sequence-specific alphaglobin mRNP complex which is associated with alpha-globin mRNA stability. This multiexon structural mRNA is thought to be retrotransposed to generate PCBP-1, an intronless gene with functions similar to that of PCBP2. This gene and PCBP-1 have paralogous genes (PCBP3 and PCBP4) which are thought to have arisen as a result of duplication events of entire genes. This gene also has two processed pseudogenes (PCBP2P1 and PCBP2P2). Multiple transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB 1:1000 - 1:2000

ΙP 0.5µg-4µg antibody for 200µg-400µg extracts

of whole cells

ELISA Recommended starting

concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Contact

€ www.abclonal.com

Immunogen Information

Gene ID Swiss Prot 5094 Q15366

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 150-250 of human hnRNP E2/PCBP2 (NP_001122383.1).

Svnonvms

HNRPE2; HNRNPE2; hnRNP-E2; hnRNP E2/PCBP2

Product Information


Source **Purification Isotype** Rabbit Affinity purification IgG

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates, using hnRNP E2/PCBP2 (A23987) at 1:2000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000


dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.



Immunoprecipitation analysis of 300ug extracts of Hela cells using 3ug hnRNP E2/PCBP2 Rabbit mAb (A23987). Western blot was performed from the immunoprecipitate using hnRNP E2/PCBP2 Rabbit mAb (A23987) at a dilition of 1:1000.