# ABclonal www.abclonal.com

# **PPBP Rabbit pAb**

Catalog No.: A24514

# **Basic Information**

#### **Observed MW**

11kDa/14kDa

#### **Calculated MW**

13kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB, ELISA

# **Cross-Reactivity**

Human

# **Background**

The protein encoded by this gene is a platelet-derived growth factor that belongs to the CXC chemokine family. This growth factor is a potent chemoattractant and activator of neutrophils. It has been shown to stimulate various cellular processes including DNA synthesis, mitosis, glycolysis, intracellular cAMP accumulation, prostaglandin E2 secretion, and synthesis of hyaluronic acid and sulfated glycosaminoglycan. It also stimulates the formation and secretion of plasminogen activator by synovial cells. The protein also is an antimicrobial protein with bactericidal and antifungal activity.

# **Recommended Dilutions**

WB

1:500 - 1:1000

# **Immunogen Information**

Gene ID 5473 Swiss Prot P02775

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 59-128 of human PPBP (NP\_002695.1).

### **Synonyms**

PPBP; B-TG1; Beta-TG; CTAP-III; CTAP3; CTAPIII; CXCL7; LA-PF4; LDGF; MDGF; NAP-2; PBP; SCYB7; TC1; TC2; TGB; TGB1; THBGB; THBGB1; platelet basic protein

# **Contact**

•

www.abclonal.com

# **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of lysates from wild type (WT) and 293T cells transfected with PPBP, using PPBP Rabbit pAb (A24514) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25ug per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.