NAGS Rabbit pAb

Catalog No.: A24553



Basic Information

Observed MW 51kDa

Calculated MW 58kDa

Category Polyclonal Antibody

Applications WB,ELISA

Cross-Reactivity Human, Mouse

Background

The N-acetylglutamate synthase gene encodes a mitochondrial enzyme that catalyzes the formation of N-acetylglutamate (NAG) from glutamate and acetyl coenzyme-A. NAG is a cofactor of carbamyl phosphate synthetase I (CPSI), the first enzyme of the urea cycle in mammals. This gene may regulate ureagenesis by altering NAG availability and, thereby, CPSI activity. Deficiencies in N-acetylglutamate synthase have been associated with hyperammonemia.

Immunogen Information

WB

Gene ID 162417

Swiss Prot Q8N159

Immunogen Recombinant protein of human NAGS.

Synonyms NAGS; AGAS; ARGA; N-acetylglutamate synthase

Contact

Product Information

www.abclonal.com

Recommended Dilutions


1:500 - 1:2000

Source Rabbit **Isotype** IgG **Purification** Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

Validation Data



Western blot analysis of lysates from wild type (WT) and 293T cells transfected with NAGS Rabbit pAb, using NAGS Rabbit pAb (A24553) at 1:2000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.