A2479

Leader in Biomolecular Solutions for Life Science

www.abclonal.com

SOX9 Rabbit pAb

Catalog No.: A2479 13 Publications

Basic Information

Observed MW
70kDa

Calculated MW
56kDa

Category
Polyclonal Antibody

Applications
WB,IHC-P,ELISA

Cross-Reactivity
Human,Mouse,Rat

Background

The protein encoded by this gene recognizes the sequence CCTTGAG along with other members of the HMG-box class DNA-binding proteins. It acts during chondrocyte differentiation and, with steroidogenic factor 1, regulates transcription of the anti-Muellerian hormone (AMH) gene. Deficiencies lead to the skeletal malformation syndrome campomelic dysplasia, frequently with sex reversal.

Recommended Dilutions

WB	1:500 - 1:2000
IHC-P	1:50 - 1:200
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
6662

Swiss Prot
P48436

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms
CMD1; SRA1; CMPD1; SRXX2; SRXY10; SOX9

Contact

www.abclonal.com

Product Information


Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20℃


Validation Data




Western blot analysis of various lysates using SOX9 Rabbit pAb (A2479) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 120s.



Immunohistochemistry analysis of paraffin-embedded Human thyroid cancer using SOX9 Rabbit pAb (A2479) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human colon using SOX9 Rabbit pAb (A2479) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse pancreas using SOX9 Rabbit pAb (A2479) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.

For research use only. Not for therapeutic or diagnostic purposes.
Please visit https://abclonal.com for a complete listing of recommended products.