A26297

Leader in Biomolecular Solutions for Life Science

www.abclonal.com

CARD8 Rabbit mAb

Catalog No.: A26297 Recombinant 1 Publications

Basic Information

Observed MW
45 kDa

Calculated MW
61 kDa

Category
SMab Recombinant Monoclonal Antibody

Applications
WB,ELISA

Cross-Reactivity
Human,Rat

Background

The protein encoded by this gene belongs to the caspase recruitment domain (CARD)-containing family of proteins, which are involved in pathways leading to activation of caspases or nuclear factor kappa-B (NFKB). This protein may be a component of the inflammasome, a protein complex that plays a role in the activation of proinflammatory caspases. It is thought that this protein acts as an adaptor molecule that negatively regulates NFKB activation, CASP1-dependent IL1B secretion, and apoptosis. Polymorphisms in this gene may be associated with a susceptibility to rheumatoid arthritis. Alternatively spliced transcript variants have been described for this gene.

Recommended Dilutions

WB	1:1000 - 1:6000
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
22900

Swiss Prot
Q9Y2G2

Immunogen
Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms
NDPP; DACAR; DAKAR; NDPP1; TUCAN; CARDINAL

Contact

www.abclonal.com

Product Information


Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20℃

Validation Data



Western blot analysis of lysates from Rat testis using CARD8 Rabbit mAb (A26297) at 1:3000 dilution incubated overnight at 4℃.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 20 s.



Western blot analysis of lysates from HUVEC cells using CARD8 Rabbit mAb (A26297) at 1:3000 dilution incubated overnight at 4℃.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 60 s.