A26879

Leader in Biomolecular Solutions for Life Science

www.abclonal.com

# CDKN2A/p19ARF Rabbit mAb

**Catalog No.: A26879** Recombinant 2 Publications

## Basic Information

**Observed MW**
21kDa

**Calculated MW**
19KDa

**Category**
Monoclonal Antibody

**Applications**
WB,ELISA

**Cross-Reactivity**
Mouse

## Background

Enables MDM2/MDM4 family protein binding activity; enzyme inhibitor activity; and protein N-terminus binding activity. Involved in several processes, including negative regulation of lymphocyte proliferation; positive regulation of apoptotic process; and regulation of transcription, DNA-templated. Acts upstream of or within several processes, including negative regulation of cellular protein metabolic process; negative regulation of mammary gland epithelial cell proliferation; and positive regulation of apoptotic process involved in mammary gland involution. Located in cytoplasm and granular component. Part of protein-containing complex. Is expressed in several structures, including bone marrow; central nervous system; dorsal root ganglion; eye; and testis. Human ortholog(s) of this gene implicated in several diseases, including breast cancer (multiple); carcinoma (multiple); hematologic cancer (multiple); melanoma (multiple); and nervous system cancer (multiple). Orthologous to human CDKN2A (cyclin dependent kinase inhibitor 2A).

## Recommended Dilutions

| | |
|---|---|
| **WB** | 1:1000 - 1:10000 |
| **ELISA** | Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements. |

## Immunogen Information

**Gene ID**
12578

**Swiss Prot**
Q64364

**Immunogen**
Synthetic peptide. This information is considered to be commercially sensitive.

**Synonyms**
Arf; p16; MTS1; Pctr1; p19ARF; p16INK4a; p19<ARF>; ARF-INK4a; INK4a-ARF; Ink4a/Arf; p16(INK4a); CDKN2A/p19ARF

## Contact

www.abclonal.com

## Product Information


**Source**
Rabbit

**Isotype**
IgG

**Purification**
Affinity purification

**Storage**
Store at -20℃

## Validation Data




Western blot analysis of various lysates using CDKN2A/p19ARF Rabbit mAb (A26879)at 1:1000 dilution incubated overnight at 4℃.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Negative control (NC): NIH/3T3
Exposure time: 0.5s.


For research use only. Not for therapeutic or diagnostic purposes.
Please visit https://abclonal.com for a complete listing of recommended products.