CYP7A1 Rabbit pAb

Catalog No.: A27766



Basic Information

Observed MW

Refer to figures

Calculated MW

58kDa

Category

Primary Antibody

Applications

IF/ICC,ELISA

Cross-Reactivity

Human

Background

This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This endoplasmic reticulum membrane protein catalyzes the first reaction in the cholesterol catabolic pathway in the liver, which converts cholesterol to bile acids. This reaction is the rate limiting step and the major site of regulation of bile acid synthesis, which is the primary mechanism for the removal of cholesterol from the body. Polymorphisms in the promoter of this gene are associated with defects in bile acid synthesis.

Recommended Dilutions

IF/ICC 1:50 - 1:200

ELISA

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific

assay requirements.

Immunogen Information

Gene ID Swiss Prot 1581 P22680

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human CYP7A1 (NP_000771.2).

Synonyms

CP7A; CYP7; CYPVII

Contact

www.abclonal.com


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.

Validation Data



Immunofluorescence analysis of Human liver tissue using CYP7A1 Rabbit pAb (A27766) at a dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L)(AS007) at 1:500 dilution. Blue: DAPI for nuclear staining. High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining.