

A27818

Leader in Biomolecular Solutions for Life Science



Fibrillarin Rabbit mAb

Catalog No.: A27818

Recombinant

Basic Information

Observed MW

37KD

Calculated MW

34kDa

Category

SMab Recombinant Monoclonal
Antibody

Applications

WB,IHC-P,IF/ICC,IP,ELISA

Cross-Reactivity

Human,Mouse,Rat

Background

This gene product is a component of a nucleolar small nuclear ribonucleoprotein (snRNP) particle thought to participate in the first step in processing preribosomal RNA. It is associated with the U3, U8, and U13 small nuclear RNAs and is located in the dense fibrillar component (DFC) of the nucleolus. The encoded protein contains an N-terminal repetitive domain that is rich in glycine and arginine residues, like fibrillarins in other species. Its central region resembles an RNA-binding domain and contains an RNP consensus sequence. Antisera from approximately 8% of humans with the autoimmune disease scleroderma recognize fibrillarin.

Recommended Dilutions

WB 1:25000 - 1:100000

IP 0.5µg-4µg antibody for
200µg-400µg extracts
of whole cells

IF/ICC 1:200 - 1:400

IHC-P 1:1000 - 1:4000

ELISA Recommended starting
concentration is 1
µg/mL. Please optimize
the concentration
based on your specific
assay requirements.

Immunogen Information

Gene ID

2091

Swiss Prot

P22087

Immunogen

A synthetic peptide corresponding to a sequence with in amino acids 111-210 of human Fibrillarin/U3 RNP (NP_001427.2).

Synonyms

FIB; FLRN; Nop1; RNU3IP1

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.09% Sodium azide,0.05% BSA,50% glycerol,pH7.3.

Contact




www.abclonal.com


Validation Data




Western blot analysis of various lysates using Fibrillarin Rabbit mAb (A27818) at 1:25000 dilution incubated overnight at 4°C.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25 µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 20s.




Immunohistochemistry analysis of paraffin-embedded Mouse liver tissue using Fibrillarin Rabbit mAb (A27818) at a dilution of 1:3000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human tonsil tissue using Fibrillarin Rabbit mAb (A27818) at a dilution of 1:3000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat skin tissue using Fibrillarin Rabbit mAb (A27818) at a dilution of 1:3000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Confocal imaging of NIH/3T3 cells using Fibrillarin Rabbit mAb (A27818, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear




Confocal imaging of HeLa cells using Fibrillarin Rabbit mAb (A27818, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear

Validation Data

staining (Blue). Objective: 100x.

staining (Blue). Objective: 100x.



Immunoprecipitation of Fibrillarin from 300 µg extracts of 293T cells was performed using 0.5 µg of Fibrillarin Rabbit mAb (A27818). Rabbit IgG isotype control (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with 1X reducing Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using Fibrillarin Rabbit mAb (A27818) at a dilution of 1:5000.