# Lipin 1 Rabbit mAb

Catalog No.: A3326 Recombinant 1 Publications



#### **Basic Information**

**Observed MW** 130kDa

**Calculated MW** 99kDa

Category SMab Recombinant Monoclonal Antibody

Applications WB, ELISA

**Cross-Reactivity** Human, Mouse, Rat

**CloneNo number** ARC1945

### **Recommended Dilutions**

WB	1:1000 - 1:2000
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## Background

This gene encodes a magnesium-ion-dependent phosphatidic acid phosphohydrolase enzyme that catalyzes the penultimate step in triglyceride synthesis including the dephosphorylation of phosphatidic acid to yield diacylglycerol. Expression of this gene is required for adipocyte differentiation and it also functions as a nuclear transcriptional coactivator with some peroxisome proliferator-activated receptors to modulate expression of other genes involved in lipid metabolism. Mutations in this gene are associated with metabolic syndrome, type 2 diabetes, acute recurrent rhabdomyolysis, and autosomal recessive acute recurrent myoglobinuria (ARARM). This gene is also a candidate for several human lipodystrophy syndromes.

# **Immunogen Information**

Gene ID 23175

Swiss Prot Q14693

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 101-200 of human Lipin 1 (Q14693).

#### **Synonyms**

PAP1; Lipin 1

# Contact

# **Product Information**

Ð www.abclonal.com

Isotype IgG


Purification Affinity purification

#### Storage

Source

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.



Western blot analysis of various lysates using Lipin 1 Rabbit mAb (A3326) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.