

EAAT2/SLC1A2 Rabbit mAb

Catalog No.: A3679 Recombinant 1 Publications

Basic Information

Observed MW 70kDa

Calculated MW 62kDa

Category

SMab Recombinant Monoclonal Antibody

Applications WB, ELISA

Cross-Reactivity Human, Mouse, Rat

CloneNo number ARC0821

Recommended Dilutions

WB	1:500 - 1:2000
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Background

This gene encodes a member of a family of solute transporter proteins. The membranebound protein is the principal transporter that clears the excitatory neurotransmitter glutamate from the extracellular space at synapses in the central nervous system. Glutamate clearance is necessary for proper synaptic activation and to prevent neuronal damage from excessive activation of glutamate receptors. Improper regulation of this gene is thought to be associated with several neurological disorders. Alternatively spliced transcript variants of this gene have been identified.

Immunogen Information

Gene ID 6506

Swiss Prot P43004

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 150-250 of human EAAT2/SLC1A2 (P43004).

Synonyms

GLT1; HBGT; DEE41; EAAT2; GLT-1; EIEE41; EAAT2/SLC1A2

Contact

Product Information

G www.abclonal.com

Source Rabbit


Isotype IgG

Purification Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of various lysates using EAAT2/SLC1A2 Rabbit mAb (A3679) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.