

A3719

Leader in Biomolecular Solutions for Life Science



[KO Validated] ACLY Rabbit mAb

Catalog No.: A3719

KO Validated

Recombinant

3 Publications

Basic Information

Observed MW

121 kDa

Calculated MW

121 kDa

Category

Monoclonal Antibody

Applications

WB,IF/ICC,IP,ELISA

Cross-Reactivity

Human,Mouse,Rat

CloneNo number

ARC0281

Recommended Dilutions

WB 1:6000 - 1:20000

IP 0.5µg-4µg antibody for
200µg-400µg extracts
of whole cells

IF/ICC 1:50 - 1:200

ELISA Recommended starting
concentration is 1
µg/mL. Please optimize
the concentration
based on your specific
assay requirements.

Contact

 www.abclonal.com

Background

ATP citrate lyase is the primary enzyme responsible for the synthesis of cytosolic acetyl-CoA in many tissues. The enzyme is a tetramer (relative molecular weight approximately 440,000) of apparently identical subunits. It catalyzes the formation of acetyl-CoA and oxaloacetate from citrate and CoA with a concomitant hydrolysis of ATP to ADP and phosphate. The product, acetyl-CoA, serves several important biosynthetic pathways, including lipogenesis and cholesterologenesis. In nervous tissue, ATP citrate-lyase may be involved in the biosynthesis of acetylcholine. Multiple transcript variants encoding distinct isoforms have been identified for this gene.

Immunogen Information

Gene ID

47

Swiss Prot

P53396

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

ACL; ATPCL; CLATP; ACLY

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of lysates from wild type (WT) and ACLY knockout (KO) 293T cells using [KO Validated] ACLY Rabbit mAb (A3719) at 1:10000 dilution incubated overnight at 4°C.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 20 s.



Western blot analysis of various lysates using [KO Validated] ACLY Rabbit mAb (A3719) at 1:10000 dilution incubated overnight at 4°C.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 20 s.




Confocal imaging of NIH/3T3 cells using [KO Validated] ACLY Rabbit mAb (A3719,dilution 1:100)(Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012,dilution 1:400) (Green). DAPI was used for nuclear staining (blue). Objective: 100x.



Confocal imaging of U-2 OS cells using [KO Validated] ACLY Rabbit mAb (A3719,dilution 1:100)(Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012,dilution 1:400) (Green). DAPI was used for nuclear staining (blue). Objective: 100x.

Validation Data



Immunoprecipitation analysis of 300 μ g extracts from Hep G2 cells using 3 μ g [KO Validated] ACLY Rabbit mAb (A3719). Western blot was performed from the immunoprecipitate using [KO Validated] ACLY Rabbit mAb (A3719) at a dilution of 1:1000.