# PTPN7 Rabbit pAb

Catalog No.: A4062



## **Basic Information**

## **Observed MW**

40kDa

#### **Calculated MW**

41kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB, ELISA

## **Cross-Reactivity**

Human

# **Background**

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This gene is preferentially expressed in a variety of hematopoietic cells, and is an early response gene in lymphokine stimulated cells. The non-catalytic N-terminus of this PTP can interact with MAP kinases and suppress the MAP kinase activities. This PTP was shown to be involved in the regulation of T cell antigen receptor (TCR) signaling, which was thought to function through dephosphorylating the molecules related to MAP kinase pathway. Multiple alternatively spliced transcript variants have been found for this gene.

## **Recommended Dilutions**

WB

1:500 - 1:2000

## **Immunogen Information**

Gene ID 5778 **Swiss Prot** 

P35236

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-190 of human PTPN7 (NP\_542155.1).

### **Synonyms**

LPTP; HEPTP; PTPNI; BPTP-4; LC-PTP; PTPN7

## **Contact**

•

www.abclonal.com

## **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of extracts of various cell lines, using PTPN7 antibody (A4062) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.