RPL3 Rabbit pAb

Catalog No.: A4082



Basic Information

Observed MW 50kDa

Calculated MW 46kDa

Category Polyclonal Antibody

Applications WB

Cross-Reactivity Human, Mouse

Background

Ribosomes, the complexes that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L3P family of ribosomal proteins and it is located in the cytoplasm. The protein can bind to the HIV-1 TAR mRNA, and it has been suggested that the protein contributes to tat-mediated transactivation. This gene is co-transcribed with several small nucleolar RNA genes, which are located in several of this gene's introns. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.

Recommended Dilutions

Immunogen Information

WB

1:200 - 1:2000

Gene ID 6122 Swiss Prot P39023

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

L3; uL3; ASC-1; TARBP-B; RPL3

Product Information


www.abclonal.com

Source Rabbit **Isotype** IgG Purification Affinity purification

Storage

Store at 4°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,pH7.3.

Validation Data



Western blot analysis of various lysates using RPL3 Rabbit pAb (A4082). Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.