Leader in Biomolecular Solutions for Life Science

# **NEDD4 Rabbit mAb**

Catalog No.: A4385 Recombinant 2 Publications



### **Basic Information**

#### **Observed MW**

140kDa

#### **Calculated MW**

149kDa

### Category

SMab Recombinant Monoclonal Antibody

### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human

#### CloneNo number

ARC0992

# **Background**

This gene is the founding member of the NEDD4 family of HECT ubiquitin ligases that function in the ubiquitin proteasome system of protein degradation. The encoded protein contains an N-terminal calcium and phospholipid binding C2 domain followed by multiple tryptophan-rich WW domains and, a C-terminal HECT ubiquitin ligase catalytic domain. It plays critical role in the regulation of a number of membrane receptors, endocytic machinery components and the tumor suppressor PTEN.

# **Recommended Dilutions**

WB

1:500 - 1:1000

# **Immunogen Information**

Gene ID 4734 Swiss Prot P46934

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 1200-1300 of human NEDD4 (P46934).

# **Synonyms**

RPF1; NEDD4-1; NEDD4

### **Contact**

•

www.abclonal.com

## **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

# **Validation Data**



Western blot analysis of extracts of various cell lines, using NEDD4 Rabbit mAb (A4385) at 1:1000

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.