ABclonal www.abclonal.com

GluR4/GluA4/GRIA4 Rabbit mAb

Catalog No.: A4593 Recombinant

Basic Information

Observed MW

101kDa

Calculated MW

101kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Mouse,Rat

CloneNo number

ARC1045

Background

Glutamate receptors are the predominant excitatory neurotransmitter receptors in the mammalian brain and are activated in a variety of normal neurophysiologic processes. These receptors are heteromeric protein complexes composed of multiple subunits, arranged to form ligand-gated ion channels. The classification of glutamate receptors is based on their activation by different pharmacologic agonists. The subunit encoded by this gene belongs to a family of AMPA (alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate)-sensitive glutamate receptors, and is subject to RNA editing (AGA->GGA; R->G). Alternative splicing of this gene results in transcript variants encoding different isoforms, which may vary in their signal transduction properties. Some haplotypes of this gene show a positive association with schizophrenia.

Recommended Dilutions

WB

1:500 - 1:2000

Immunogen Information

Gene ID 2893 Swiss Prot P48058

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 800-902 of human GluR4/GluA4/GRIA4 (P48058).

Synonyms

GLUR4; GLURD; GluA4; GLUR4C; NEDSGA; GluA4-ATD; GluR4/GluA4/GRIA4

Contact

•

www.abclonal.com

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of extracts of various cell lines, using GluR4/GluA4/GRIA4Rabbit mAb (A4593) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.