ABclomal www.abclonal.com

ATGL/PNPLA2 Rabbit mAb

Catalog No.: A5126 Recombinant 7 Publications

Basic Information

Observed MW

55kDa

Calculated MW

55kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC1192

Background

This gene encodes an enzyme which catalyzes the first step in the hydrolysis of triglycerides in adipose tissue. Mutations in this gene are associated with neutral lipid storage disease with myopathy.

Recommended Dilutions

WB 1:1000 - 1:2000

ELISA

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID Swiss Prot 57104 Q96AD5

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

ATGL; TTS2; PEDF-R; FP17548; TTS-2.2; iPLA2zeta; 1110001C14Rik; ATGL/PNPLA2

Contact

0 www.abclonal.com

Product Information


Source **Isotype Purification** Rabbit Affinity purification IgG

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of various lysates using ATGL/PNPLA2 Rabbit mAb (A5126) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.