

A5271SP

Leader in Biomolecular Solutions for Life Science



[KO Validated] CD63 Rabbit pAb

Catalog No.: A5271SP

KO Validated

78 Publications

Basic Information

Observed MW

30-65 kDa

Calculated MW

26 kDa/23 kDa/17 kDa

Category

Polyclonal Antibody

Applications

WB,IHC-P,ELISA

Cross-Reactivity

Human

Background

The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. The encoded protein is a cell surface glycoprotein that is known to complex with integrins. It may function as a blood platelet activation marker. Deficiency of this protein is associated with Hermansky-Pudlak syndrome. Also this gene has been associated with tumor progression. Alternative splicing results in multiple transcript variants encoding different protein isoforms.

Recommended Dilutions

WB 1:1000 - 1:2000

IHC-P 1:2000 - 1:8000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. For high-ratio antibody dilutions (≥1:10000), a sequential dilution method is strongly recommended to ensure measurement accuracy.

Immunogen Information

Gene ID

967

Swiss Prot

P08962

Immunogen

This information is considered to be commercially sensitive.

Synonyms

MLA1; ME491; LAMP-3; OMA81H; TSPAN30; CD63

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS, pH 7.3, containing 50% glycerol. Preserved with Proclin300 or sodium azide. May contain 0.05% BSA as specified on the Certificate of Analysis.

Contact



www.abclonal.com

Validation Data



Western blot analysis of lysates from wild type (WT) and CD63 knockout (KO) 293T cells using [KO Validated] CD63 Rabbit pAb (A5271SP) at 1:1000 dilution incubated overnight at 4°C.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90 s.



Immunohistochemistry analysis of paraffin-embedded Human malignant melanoma tissue using [KO Validated] CD63 Rabbit pAb (A5271SP) at a dilution of 1:8000 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human tonsil tissue using [KO Validated] CD63 Rabbit pAb (A5271SP) at a dilution of 1:8000 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.