

A5865

Leader in Biomolecular Solutions for Life Science



# CD71/Transferrin Receptor Rabbit pAb

Catalog No.: A5865

63 Publications

## Basic Information

### Observed MW

100kDa

### Calculated MW

85kDa

### Category

Polyclonal Antibody

### Applications

WB,IHC-P,IF/ICC,ELISA

### Cross-Reactivity

Human,Mouse,Rat

## Background

This gene encodes a cell surface receptor necessary for cellular iron uptake by the process of receptor-mediated endocytosis. This receptor is required for erythropoiesis and neurologic development. Multiple alternatively spliced variants have been identified.

## Recommended Dilutions

**WB** 1:1500 - 1:9000

**IHC-P** 1:50 - 1:200

**IF/ICC** 1:50 - 1:200

**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID

7037

### Swiss Prot

P02786

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-140 of human CD71/Transferrin Receptor (NP\_001121620.1).

### Synonyms

T9; TR; TFR; p90; CD71; TFR1; TRFR; IMD46; CD71/Transferrin Receptor

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.

## Contact



[www.abclonal.com](http://www.abclonal.com)

## Validation Data



Western blot analysis of lysates from HeLa cells using CD71/Transferrin Receptor Rabbit pAb (A5865) at 1:3000 dilution incubated overnight at 4°C.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 10s.




Immunohistochemistry analysis of paraffin-embedded Rat kidney using CD71/Transferrin Receptor Rabbit pAb (A5865) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human liver cancer using CD71/Transferrin Receptor Rabbit pAb (A5865) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma using CD71/Transferrin Receptor Rabbit pAb (A5865) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human placenta using CD71/Transferrin Receptor Rabbit pAb (A5865) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunofluorescence analysis of PC-12 cells using CD71/Transferrin Receptor Rabbit pAb (A5865) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.