A6614

Leader in Biomolecular Solutions for Life Science

www.abclonal.com

[KO Validated] Histone H2A.Z Rabbit pAb

Catalog No.: A6614 KO Validated 2 Publications

Basic Information

Observed MW
15 kDa

Calculated MW
13 kDa

Category
Polyclonal Antibody

Applications
WB,ELISA

Cross-Reactivity
Human,Mouse,Rat

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene encodes a replication-independent member of the histone H2A family that is distinct from other members of the family. Studies in mice have shown that this particular histone is required for embryonic development and indicate that lack of functional histone H2A leads to embryonic lethality.

Recommended Dilutions

WB	1:500 - 1:5000
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
3015

Swiss Prot
P0C0S5

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms
H2AZ; H2A.z; H2A/z; H2AFZ; H2A.Z-1; .Z

Contact

www.abclonal.com

Product Information


Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20℃

Validation Data



Western blot analysis of various lysates using [KO Validated] Histone H2A.Z Rabbit pAb (A6614) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.



Western blot analysis of lysates from wild type (WT) and Histone H2A.Z knockout (KO) 293T cells, using [KO Validated] Histone H2A.Z Rabbit pAb (A6614) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 1s.