# ABclonal www.abclonal.com

## INPP5J Rabbit pAb

Catalog No.: A6631

## **Basic Information**

#### **Observed MW**

107kDa

#### **Calculated MW**

107kDa

#### Category

Polyclonal Antibody

## **Applications**

WB,IF/ICC,ELISA

## **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

Predicted to enable phosphatidylinositol-3,4,5-trisphosphate 5-phosphatase activity and phosphatidylinositol-4,5-bisphosphate 5-phosphatase activity. Predicted to be involved in inositol phosphate dephosphorylation; negative regulation of peptidylserine phosphorylation; and phosphatidylinositol dephosphorylation. Predicted to act upstream of or within negative regulation of neuron projection development. Located in cytoplasm and ruffle.

## **Recommended Dilutions**

**WB** 1:500 - 1:2000

**IF/ICC** 1:10 - 1:100

**ELISA** Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific

assay requirements.

## **Immunogen Information**

**Gene ID**27124

Swiss Prot
Q15735

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

#### **Synonyms**

PIPP; INPP5; PIB5PA; INPP5J

## **Contact**

www.abclonal.com

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of various lysates using INPP5J Rabbit pAb (A6631) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000


dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.



Immunofluorescence analysis of U2OS cells using INPP5J Rabbit pAb (A6631). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.