

A7042

Leader in Biomolecular Solutions for Life Science



# NMT2 Rabbit pAb

Catalog No.: A7042

## Basic Information

### Observed MW

57kDa

### Calculated MW

57kDa

### Category

Polyclonal Antibody

### Applications

WB,IF/ICC,ELISA

### Cross-Reactivity

Human,Mouse

## Background

This gene encodes one of two N-myristoyltransferase proteins. N-terminal myristoylation is a lipid modification that is involved in regulating the function and localization of signaling proteins. The encoded protein catalyzes the addition of a myristoyl group to the N-terminal glycine residue of many signaling proteins, including the human immunodeficiency virus type 1 (HIV-1) proteins, Gag and Nef. Alternative splicing results in multiple transcript variants.

## Recommended Dilutions

WB 1:500 - 1:2000

IF/ICC 1:50 - 1:100

## Immunogen Information

### Gene ID

9397

### Swiss Prot

O60551

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-120 of human NMT2 (NP\_004799.1).

### Synonyms

NMT2

## Contact



[www.abclonal.com](http://www.abclonal.com)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of extracts of various cell lines, using NMT2 antibody (A7042) at 1:1000 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.



Immunofluorescence analysis of A-549 cells using NMT2 antibody (A7042). Blue: DAPI for nuclear staining.