LDLRAP1 Rabbit pAb

Catalog No.: A7093



Basic Information

Observed MW 36kDa

Calculated MW 34kDa

Category **Polyclonal Antibody**

Applications WB, IF/ICC, ELISA

Cross-Reactivity Human, Mouse, Rat

Background

The protein encoded by this gene is a cytosolic protein which contains a phosphotyrosine binding (PTD) domain. The PTD domain has been found to interact with the cytoplasmic tail of the LDL receptor. Mutations in this gene lead to LDL receptor malfunction and cause the disorder autosomal recessive hypercholesterolaemia.

Recommended Dilutions

WB	1:500 - 1:1000
IF/ICC	1:50 - 1:100
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID	Swiss Prot
26119	Q5SW96
T	

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 59-308 of human LDLRAP1 (NP_056442.2).

Synonyms

ARH; ARH1; ARH2; FHCB1; FHCB2; FHCL4; LDLRAP1

Contact

Product Information

Ð www.abclonal.com

Source Rabbit


Isotype IgG

Purification Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates using LDLRAP1 Rabbit pAb (A7093) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.



Immunofluorescence analysis of A-549 cells using LDLRAP1 Rabbit pAb (A7093).Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution.