# ABclonal www.abclonal.com

# NFIL3 Rabbit pAb

Catalog No.: A7294

# **Basic Information**

#### **Observed MW**

60kDa

#### **Calculated MW**

51kDa

#### Category

Polyclonal Antibody

### **Applications**

WB,IF/ICC,ELISA

### **Cross-Reactivity**

Human

# **Background**

The protein encoded by this gene is a transcriptional regulator that binds as a homodimer to activating transcription factor (ATF) sites in many cellular and viral promoters. The encoded protein represses PER1 and PER2 expression and therefore plays a role in the regulation of circadian rhythm. Three transcript variants encoding the same protein have been found for this gene.

# **Recommended Dilutions**

**WB** 1:500 - 1:2000

**IF/ICC** 1:50 - 1:100

**ELISA** Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID**4783

Swiss Prot
Q16649

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 133-462 of human NFIL3 (NP\_005375.2).

# **Synonyms**

E4BP4; IL3BP1; NFIL3A; NF-IL3A; NFIL3

### **Contact**

www.abclonal.com


# **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

# Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of lysates from SW620 cells, using NFIL3 Rabbit pAb (A7294) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.