Leader in Biomolecular Solutions for Life Science

# NAA40 Rabbit pAb

Catalog No.: A7385



### **Basic Information**

# **Observed MW**

27kDa

### **Calculated MW**

27kDa

#### Category

Polyclonal Antibody

# **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human

# **Background**

Enables H2A histone acetyltransferase activity; H4 histone acetyltransferase activity; and peptide-serine-N-acetyltransferase activity. Involved in N-terminal protein amino acid acetylation; histone H2A acetylation; and histone H4 acetylation. Located in centriolar satellite; cytosol; and nucleoplasm.

# **Recommended Dilutions**

WB

1:500 - 1:1000

# **Immunogen Information**

**Gene ID** 79829

Swiss Prot Q86UY6

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-237 of human NAA40 (NP\_079047.2).

### **Synonyms**

NatD; NAT11; PATT1; hNatD; NAA40

# **Contact**

•

www.abclonal.com

# **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

# Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of lysates from 293T cells, using NAA40 Rabbit pAb (A7385) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.