

A7463

Leader in Biomolecular Solutions for Life Science



# DAAM2 Rabbit pAb

Catalog No.: A7463

1 Publications

## Basic Information

### Observed MW

123kDa

### Calculated MW

123kDa

### Category

Polyclonal Antibody

### Applications

WB,IF/ICC,ELISA

### Cross-Reactivity

Human,Mouse,Rat

## Background

Predicted to enable actin binding activity and small GTPase binding activity. Predicted to be involved in nervous system development and regulation of Wnt signaling pathway. Predicted to act upstream of or within determination of left/right symmetry. Located in extracellular exosome.

## Recommended Dilutions

WB	1:500 - 1:2000
IF/ICC	1:50 - 1:200

## Immunogen Information

<b>Gene ID</b>	<b>Swiss Prot</b>
23500	Q86T65

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-86 of human DAAM2 (NP\_056160.2).

### Synonyms

NPHS24; dJ90A20A.1; DAAM2

## Contact

 [www.abclonal.com](http://www.abclonal.com)

## Product Information


<b>Source</b>	<b>Isotype</b>	<b>Purification</b>
Rabbit	IgG	Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of various lysates using DAAM2 Rabbit pAb (A7463) at 1:1000 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 90s.



Immunofluorescence analysis of U2OS cells using DAAM2 Rabbit pAb (A7463) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of L929 cells using DAAM2 Rabbit pAb (A7463) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of H9C2 cells using DAAM2 Rabbit pAb (A7463) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.