# FGFR4 Rabbit pAb

Catalog No.: A7555 1 Publications



#### **Basic Information**

**Observed MW** 110kDa

**Calculated MW** 88kDa

Category **Polyclonal Antibody** 

Applications WB, ELISA

**Cross-Reactivity** Human, Mouse, Rat

# Background

The protein encoded by this gene is a tyrosine kinase and cell surface receptor for fibroblast growth factors. The encoded protein is involved in the regulation of several pathways, including cell proliferation, cell differentiation, cell migration, lipid metabolism, bile acid biosynthesis, vitamin D metabolism, glucose uptake, and phosphate homeostasis. This protein consists of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment, and a cytoplasmic tyrosine kinase domain. The extracellular portion interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation.

## **Recommended Dilutions**

WB	1:500 - 1:2000
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

Gene ID 2264

**Swiss Prot** P22455

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 22-150 of human FGFR4 (NP\_998812.1).

**Synonyms** 

TKF; JTK2; CD334; FGFR4

#### Contact

### **Product Information**

Ð www.abclonal.com

Source Rabbit


Isotype IgG

Purification Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

# Validation Data



Western blot analysis of various lysates using FGFR4 Rabbit pAb (A7555) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.