Leader in Biomolecular Solutions for Life Science

# **DAZ2 Rabbit pAb**

Catalog No.: A7600



## **Basic Information**

### **Observed MW**

63kDa

#### **Calculated MW**

63kDa

#### Category

Polyclonal Antibody

## **Applications**

WB, ELISA

#### **Cross-Reactivity**

Mouse

# **Background**

This gene is a member of the DAZ gene family and is a candidate for the human Y-chromosomal azoospermia factor (AZF). Its expression is restricted to premeiotic germ cells, particularly in spermatogonia. It encodes an RNA-binding protein that is important for spermatogenesis. Four copies of this gene are found on chromosome Y within palindromic duplications; one pair of genes is part of the P2 palindrome and the second pair is part of the P1 palindrome. Each gene contains a 2.4 kb repeat including a 72-bp exon, called the DAZ repeat; the number of DAZ repeats is variable and there are several variations in the sequence of the DAZ repeat. Each copy of the gene also contains a 10.8 kb region that may be amplified; this region includes five exons that encode an RNA recognition motif (RRM) domain. This gene contains one copy of the 10.8 kb repeat. Alternative splicing results in multiple transcript variants encoding different isoforms.

# **Recommended Dilutions**

WB

1:500 - 1:2000

# Immunogen Information

**Gene ID** 57055

**Swiss Prot** 

Q13117

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-200 of human DAZ2 (NP 001005785.1).

# **Synonyms**

pDP1678; DAZ2

### **Contact**



www.abclonal.com

## **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of extracts of mouse testis, using DAZ2 antibody (A7600) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.