

A8179

Leader in Biomolecular Solutions for Life Science



# Nuclear Matrix Protein p84 (THOC1) Rabbit pAb

Catalog No.: A8179

1 Publications

## Basic Information

### Observed MW

85kDa

### Calculated MW

76kDa

### Category

Polyclonal Antibody

### Applications

WB,IHC-P,IF/ICC,ELISA

### Cross-Reactivity

Human,Mouse,Rat

## Background

Predicted to enable DNA binding activity and RNA binding activity. Involved in several processes, including negative regulation of DNA damage checkpoint; regulation of nucleobase-containing compound metabolic process; and viral mRNA export from host cell nucleus. Located in cytoplasm and nuclear speck. Part of THO complex part of transcription export complex. Colocalizes with chromosome, telomeric region.

## Recommended Dilutions

**WB** 1:500 - 1:1000

**IHC-P** 1:50 - 1:200

**IF/ICC** 1:50 - 1:100

**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID

9984

### Swiss Prot

Q96FV9

### Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### Synonyms

P84; HPR1; P84N5; DFNA86; Nuclear Matrix Protein p84 (THOC1)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## Contact




[www.abclonal.com](http://www.abclonal.com)


## Validation Data




Western blot analysis of various lysates using Nuclear Matrix Protein p84 (THOC1) Rabbit pAb (A8179) at 1:1000 dilution.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 1s.




Western blot analysis of various lysates using Nuclear Matrix Protein p84 (THOC1) Rabbit pAb (A8179) at 1:1000 dilution.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 30s.



Immunohistochemistry analysis of paraffin-embedded Human esophageal cancer using Nuclear Matrix Protein p84 (THOC1) Rabbit pAb (A8179) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse lung using Nuclear Matrix Protein p84 (THOC1) Rabbit pAb (A8179) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat ovary using Nuclear Matrix Protein p84 (THOC1) Rabbit pAb (A8179) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.

## Validation Data

---



Immunofluorescence analysis of U2OS cells using Nuclear Matrix Protein p84 (THOC1) Rabbit pAb (A8179) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.