# ABclonal www.abclonal.com

# RNF34 Rabbit pAb

Catalog No.: A8517 1 Publications

### **Basic Information**

### **Observed MW**

42kDa

### **Calculated MW**

42kDa

### Category

Polyclonal Antibody

### **Applications**

WB,IF/ICC,ELISA

### **Cross-Reactivity**

Human

## Background

The protein encoded by this gene contains a RINF finger, a motif known to be involved in protein-protein and protein-DNA interactions. This protein interacts with DNAJA3/hTid-1, which is a DnaJ protein reported to function as a modulator of apoptosis. Overexpression of this gene in Hela cells was shown to confer the resistance to TNF-alpha induced apoptosis, suggesting an anti-apoptotic function of this protein. This protein can be cleaved by caspase-3 during the induction of apoptosis. This protein also targets p53 and phospho-p53 for degradation. Alternatively splicing results in multiple transcript variants encoding distinct isoforms.

# **Recommended Dilutions**

WB 1:500 - 1:2000

IF/ICC 1:50 - 1:200

Recommended starting **ELISA** 

> concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

### **Immunogen Information**

**Gene ID Swiss Prot** 80196 Q969K3

### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

## **Synonyms**

RFI; RIF; RIFF; hRFI; CARP1; CARP-1; RNF34

### **Contact**

www.abclonal.com €

### **Product Information**


Source **Isotype Purification** Rabbit Affinity purification IgG

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of various lysates using RNF34 Rabbit pAb (A8517) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.