ABclonal www.abclonal.com

[KO Validated] B4GALT1 Rabbit pAb

Catalog No.: A8546

KO Validated 3 Publications

Background

Basic Information

Observed MW

55kDa

Calculated MW

44kDa

Category

Polyclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Human

Recommended Dilutions

WB 1:500 - 1:2000

ELISA

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

cleaved to form the soluble lactose synthase.

Gene ID Swiss Prot 2683 P15291

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 45-210 of human B4GALT1 (NP 001488.2).

This gene is one of seven beta-1,4-galactosyltransferase (beta4GalT) genes. They

the biosynthesis of different glycoconjugates and saccharide structures. As type II

transmembrane anchor. By sequence similarity, the beta4GalTs form four groups:

encode type II membrane-bound glycoproteins that appear to have exclusive specificity for the donor substrate UDP-galactose; all transfer galactose in a beta1,4 linkage to similar acceptor sugars: GlcNAc, Glc, and Xyl. Each beta4GalT has a distinct function in

membrane proteins, they have an N-terminal hydrophobic signal sequence that directs the protein to the Golgi apparatus and which then remains uncleaved to function as a

beta4GalT1 and beta4GalT2, beta4GalT3 and beta4GalT4, beta4GalT5 and beta4GalT6, and beta4GalT7. This gene is unique among the beta4GalT genes because it encodes an

enzyme that participates both in glycoconjugate and lactose biosynthesis. For the first

activity, the enzyme adds galactose to N-acetylglucosamine residues that are either monosaccharides or the nonreducing ends of glycoprotein carbohydrate chains. The

second activity is restricted to lactating mammary tissues where the enzyme forms a

heterodimer with alpha-lactalbumin to catalyze UDP-galactose + D-glucose <=> UDP + lactose. The two enzymatic forms result from alternate transcription initiation sites and post-translational processing. Two transcripts, which differ only at the 5' end, with approximate lengths of 4.1 kb and 3.9 kb encode the same protein. The longer transcript encodes the type II membrane-bound, trans-Golgi resident protein involved in glycoconjugate biosynthesis. The shorter transcript encodes a protein which is

Synonyms

GT1; GTB; CDG2D; GGTB2; B4GAL-T1; beta4Gal-T1; T1

Contact

0 www.abclonal.com

Product Information


Source Isotype Purification Rabbit IgG Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of lysates from wild type (WT) and B4GALT1 knockout (KO) HeLa cells, using [KO Validated] B4GALT1 Rabbit pAb (A8546) at 1:3000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 $\,$

dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.