

A9359

Leader in Biomolecular Solutions for Life Science



# NPY5R Rabbit mAb

Catalog No.: A9359 **Recombinant**

## Basic Information

### Observed MW

50kDa

### Calculated MW

51kDa

### Category

SMab Recombinant Monoclonal Antibody

### Applications

WB,IF/ICC,ELISA

### Cross-Reactivity

Human,Mouse,Rat

### CloneNo number

ARC2758

## Background

The protein encoded by this gene is a receptor for neuropeptide Y and peptide YY. The encoded protein appears to be involved in regulating food intake, with defects in this gene being associated with eating disorders. Also, the encoded protein is involved in a pathway that protects neuroblastoma cells from chemotherapy-induced cell death, providing a possible therapeutic target against neuroblastoma. Three transcript variants encoding the same protein have been found for this gene.

## Recommended Dilutions

<b>WB</b>	1:500 - 1:1000
<b>IF/ICC</b>	1:50 - 1:200

## Immunogen Information

<b>Gene ID</b>	<b>Swiss Prot</b>
4889	Q15761

### Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human NPY5R (Q15761).

### Synonyms

NPYR5; NPY5-R; NPY5-R; NPY5R

## Contact

 [www.abclonal.com](http://www.abclonal.com)

## Product Information


<b>Source</b>	<b>Isotype</b>	<b>Purification</b>
Rabbit	IgG	Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.


## Validation Data



Western blot analysis of extracts of various cell lines, using (A9359) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 0.5s.



Immunofluorescence analysis of Neuro-2a cells using NPY5R Rabbit mAb (A9359) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of SH-SY5Y cells using NPY5R Rabbit mAb (A9359) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.