

Catalog No.: A9794 1 Publications



Basic Information

Observed MW Refer to Figures

Calculated MW 36kDa

Category **Polyclonal Antibody**

Applications WB, ELISA

Cross-Reactivity Arabidopsis thaliana

Background

Encodes brassinosteroid (BR) signalling protein that accumulates in the nucleus as dephosphorylated form in response to BRs. Is phosphorylated by the BIN2 GSK3 kinase. It synergistically interacts with BIM1 to bind to E box sequences (CANNTG). The protein contains a nuclear localization signal (NLS), followed by a highly conserved aminoterminal domain (N) shared by all family members, a BIN2 phosphorylation domain (P), a PEST motif, involved in protein degradation in the absence of BR, and a carboxylterminal domain. BES1 can interact with the ELF6 and REF6 Jumonji N/C-domain containing proteins and may direct them to modify histone methylation upstream of some brassinosteroid responsive-genes. Works with BRAVO to regulate QC division in the root.

Recommended Dilutions

WB	1:500 - 1:2000
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID 838518

Swiss Prot Q9LN63

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

107 PROTEIN; BRASSINAZOLE-RESISTANT 2; BRI1-EMS-SUPPRESSOR 1; BZR2; F18014.7; F18014_7

Contact

Product Information

G www.abclonal.com Source Rabbit

Isotype IgG

Purification Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.