# Rabbit anti TAP-Tag pAb

Catalog No.: AE015



## **Basic Information**

**Observed MW** Refer to Figures

Calculated MW 48kDa

**Category** Polyclonal Antibody

Applications WB,IP

Cross-Reactivity Species independent

# Background

Protein tags are peptide sequences genetically grafted onto a recombinant protein. Often these tags are removable by chemical agents or by enzymatic means, such as proteolysis or intein splicing. Tags are attached to proteins for various purposes.Epitope tags are short peptide sequences which are chosen because high-affinity antibodies can be reliably produced in many different species. These are usually derived from viral genes, which explain their high immunoreactivity. Epitope tags include V5-tag, Myc-tag, HA-tag and NE-tag. These tags are particularly useful for western blotting, immunofluorescence and immunoprecipitation experiments, although they also find use in antibody purification.

# Recommended Dilutions Immunogen Information

WB	1:1000 - 1:5000	Gene ID	Swiss Prot
IP	1:100 - 1:500	<b>Immunogen</b> Synthetic peptide. This information is considered to be commercially sensitive.	

Synonyms


TAP;TAP tag;TAP-tag

Contact		Product Information		
ଚ	www.abclonal.com	<b>Source</b> Rabbit	<b>Isotype</b> IgG	<b>Purification</b> Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of lysates from fusion proteins, using Rabbit anti TAP-Tag pAb (AE015).

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.