

AE021

Leader in Biomolecular Solutions for Life Science



Mouse anti TAP-Tag mAb

Catalog No.: AE021

1 Publications

Basic Information

Observed MW

75kDa (Recombinant TAP protein)

Calculated MW

Category

Monoclonal Antibody

Applications

WB,ELISA

Cross-Reactivity

Species independent

CloneNo number

AMC0510

Background

Protein tags are peptide sequences genetically grafted onto a recombinant protein. Often these tags are removable by chemical agents or by enzymatic means, such as proteolysis or intein splicing. Tags are attached to proteins for various purposes. Epitope tags are short peptide sequences which are chosen because high-affinity antibodies can be reliably produced in many different species. These are usually derived from viral genes, which explain their high immunoreactivity. Epitope tags include V5-tag, Myc-tag, HA-tag and NE-tag. These tags are particularly useful for western blotting, immunofluorescence and immunoprecipitation experiments, although they also find use in antibody purification.

Recommended Dilutions

WB 1:5000 - 1:10000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

Swiss Prot

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

TAP;TAP tag;TAP-tag

Contact

 www.abclonal.com

Product Information

Source

Mouse

Isotype

IgG1

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of recombinant TAP protein using mouse anti TAP-Tag mAb (AE021) at dilution of 1:5000.

Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L) (AS003) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.