mCherry Rabbit pAb

Catalog No.: AE171 1 Publications



Basic Information

Observed MW 55kDa

Calculated MW

Category Polyclonal Antibody

Applications WB,ELISA

Cross-Reactivity Species independent

Background

Protein tags are peptide sequences genetically grafted onto a recombinant protein. Often these tags are removable by chemical agents or by enzymatic means, such as proteolysis or intein splicing. Tags are attached to proteins for various purposes.Epitope tags are short peptide sequences which are chosen because high-affinity antibodies can be reliably produced in many different species. These are usually derived from viral genes, which explain their high immunoreactivity. Epitope tags include V5-tag, Myc-tag, HA-tag and NE-tag. These tags are particularly useful for western blotting, immunofluorescence and immunoprecipitation experiments, although they also find use in antibody purification.

Recommended Dilutions

WB	1:2000 - 1:10000
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene	ID	

Swiss Prot

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-236 of mCherry tag.

Synonyms

mCherry; mCherry tag; mCherry-tag

Product Information

www.abclonal.com

Isotype IgG **Purification** Affinity purification


Storage

Source

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

Validation Data



Western blot analysis of lysates from wild type (WT) and 293F cells transfected with CT55-mCherry using mCherry Rabbit pAb (AE171) at 1:10000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1s.