Phospho-VASP-S157 Rabbit pAb

Catalog No.: AP0213



Basic Information

Observed MW 50kDa

Calculated MW 40kDa

Category Polyclonal Antibody

Applications WB

Cross-Reactivity Human, Mouse, Rat

Background

Vasodilator-stimulated phosphoprotein (VASP) is a member of the Ena-VASP protein family. Ena-VASP family members contain an EHV1 N-terminal domain that binds proteins containing E/DFPPPYD/E motifs and targets Ena-VASP proteins to focal adhesions. In the mid-region of the protein, family members have a proline-rich domain that binds SH3 and WW domain-containing proteins. Their C-terminal EVH2 domain mediates tetramerization and binds both G and F actin. VASP is associated with filamentous actin formation and likely plays a widespread role in cell adhesion and motility. VASP may also be involved in the intracellular signaling pathways that regulate integrin-extracellular matrix interactions. VASP is regulated by the cyclic nucleotidedependent kinases PKA and PKG.

Recommended Dilutions

Immunogen Information

WB

1:500 - 1:1000

Swiss Prot P50552

Immunogen

Gene ID

7408

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

VASP; Phospho-VASP-S157

Contact

Product Information

www.abclonal.com

Source Rabbit **Isotype** IgG **Purification** Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Validation Data

