ABclonal www.abclonal.com

Phospho-ABL1-Y393/ABL2-Y429 Rabbit pAb

Catalog No.: AP0282

Basic Information

Observed MW

210kDa

Calculated MW

122kDa/124kDa/60kDa/114-128kDa

Category

Polyclonal Antibody

Applications

WB,IF/ICC

Cross-Reactivity

Human, Mouse

Background

This gene is a protooncogene that encodes a protein tyrosine kinase involved in a variety of cellular processes, including cell division, adhesion, differentiation, and response to stress. The activity of the protein is negatively regulated by its SH3 domain, whereby deletion of the region encoding this domain results in an oncogene. The ubiquitously expressed protein has DNA-binding activity that is regulated by CDC2-mediated phosphorylation, suggesting a cell cycle function. This gene has been found fused to a variety of translocation partner genes in various leukemias, most notably the t(9;22) translocation that results in a fusion with the 5' end of the breakpoint cluster region gene (BCR; MIM:151410). Alternative splicing of this gene results in two transcript variants, which contain alternative first exons that are spliced to the remaining common exons. [provided by RefSeq, Aug 2014]

Recommended Dilutions

WB 1:500 - 1:1000

IF/ICC 1:100 - 1:200

Immunogen Information

Gene ID Swiss Prot25/27
P00519/P42684

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

ABL1/ABL2; Phospho-ABL1-Y393/ABL2-Y429

Contact

www.abclonal.com

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Validation Data



Western blot analysis of lysates from HL60 cells using Phospho-ABL1-Y393/ABL2-Y429 Rabbit pAb (AP0282) and the same antibody preincubated with blocking peptide. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% BSA.