

AP0286

Leader in Biomolecular Solutions for Life Science



Phospho-CD117/c-Kit-Y936 Rabbit pAb

Catalog No.: AP0286

Basic Information

Observed MW

125kDa/140kDa

Calculated MW

110kDa

Category

Polyclonal Antibody

Applications

WB

Cross-Reactivity

Human,Rat

Background

This gene encodes a receptor tyrosine kinase. This gene was initially identified as a homolog of the feline sarcoma viral oncogene v-kit and is often referred to as proto-oncogene c-Kit. The canonical form of this glycosylated transmembrane protein has an N-terminal extracellular region with five immunoglobulin-like domains, a transmembrane region, and an intracellular tyrosine kinase domain at the C-terminus. Upon activation by its cytokine ligand, stem cell factor (SCF), this protein phosphorylates multiple intracellular proteins that play a role in the proliferation, differentiation, migration and apoptosis of many cell types and thereby plays an important role in hematopoiesis, stem cell maintenance, gametogenesis, melanogenesis, and in mast cell development, migration and function. This protein can be a membrane-bound or soluble protein. Mutations in this gene are associated with gastrointestinal stromal tumors, mast cell disease, acute myelogenous leukemia, and piebaldism. Multiple transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB 1:500 - 1:2000

Immunogen Information

Gene ID

3815

Swiss Prot

P10721

Immunogen

A phospho specific peptide corresponding to residues surrounding Y936 of human KIT

Synonyms

PBT; SCFR; C-Kit; CD117; MASTC; Phospho-CD117/c-Kit-Y936

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Validation Data



Western blot analysis of extracts from C6 cells using Phospho-CD117/c-Kit-Y936 antibody (AP0286).
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% BSA.