

AP0396

Leader in Biomolecular Solutions for Life Science



Phospho-MAPKAPK-2/MK2-T334 Rabbit pAb

Catalog No.: AP0396

2 Publications

Basic Information

Observed MW

Calculated MW

46kDa

Category

Polyclonal Antibody

Applications

IHC-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a member of the Ser/Thr protein kinase family. This kinase is regulated through direct phosphorylation by p38 MAP kinase. In conjunction with p38 MAP kinase, this kinase is known to be involved in many cellular processes including stress and inflammatory responses, nuclear export, gene expression regulation and cell proliferation. Heat shock protein HSP27 was shown to be one of the substrates of this kinase in vivo. Two transcript variants encoding two different isoforms have been found for this gene.

Recommended Dilutions

IHC-P 1:50 - 1:100

Immunogen Information

Gene ID

9261

Swiss Prot

P49137

Immunogen

A synthetic phosphorylated peptide around T334 of human MAPKAPK2 (NP_116584.2?).

Synonyms

MK2; MK-2; MAPKAP-K2; Phospho-MAPKAPK-2/MK2-T334

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG


Purification

Affinity purification


Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.




Immunohistochemistry analysis of paraffin-embedded Mouse kidney using Phospho-MAPKAPK-2/MK2-T334 Rabbit pAb (AP0396) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of paraffin-embedded Rat heart using Phospho-MAPKAPK-2/MK2-T334 Rabbit pAb (AP0396) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of paraffin-embedded Human colon cancer using Phospho-MAPKAPK-2/MK2-T334 Rabbit pAb (AP0396) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded Human lung cancer using Phospho-MAPKAPK-2/MK2-T334 Rabbit pAb (AP0396) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded Rat brain using Phospho-MAPKAPK-2/MK2-T334 Rabbit pAb (AP0396) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded Human placenta using Phospho-MAPKAPK-2/MK2-T334 Rabbit pAb (AP0396) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.