# Phospho-p53-T18 Rabbit pAb

Catalog No.: AP0464 2 Publications



## **Basic Information**

**Observed MW** 54kDa

**Calculated MW** 44kDa

Category **Polyclonal Antibody** 

Applications WB, ELISA

**Cross-Reactivity** Human

## Background

This gene encodes a tumor suppressor protein containing transcriptional activation, DNA binding, and oligomerization domains. The encoded protein responds to diverse cellular stresses to regulate expression of target genes, thereby inducing cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. Mutations in this gene are associated with a variety of human cancers, including hereditary cancers such as Li-Fraumeni syndrome. Alternative splicing of this gene and the use of alternate promoters result in multiple transcript variants and isoforms. Additional isoforms have also been shown to result from the use of alternate translation initiation codons from identical transcript variants (PMIDs: 12032546, 20937277).

# **Recommended Dilutions**

WB 1:500 - 1:2000 ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

Gene ID 7157

**Swiss Prot** P04637

#### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

#### **Synonyms**

P53; BCC7; LFS1; BMFS5; TRP53; Phospho-p53-T18

Contact
---------

### **Product Information**

€ www.abclonal.com

Isotype IgG

Purification Affinity purification


#### Storage

Source

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

# Validation Data



Western blot analysis of lysates from MDA cells, using Phospho-p53-T18 Rabbit pAb (AP0464).

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% BSA.