# ABclonal www.abclonal.com

# Phospho-PFKFB2-S483 Rabbit pAb

Catalog No.: AP0784

#### **Basic Information**

#### **Observed MW**

55kDa

#### **Calculated MW**

58kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human

## **Background**

The protein encoded by this gene is involved in both the synthesis and degradation of fructose-2,6-bisphosphate, a regulatory molecule that controls glycolysis in eukaryotes. The encoded protein has a 6-phosphofructo-2-kinase activity that catalyzes the synthesis of fructose-2,6-bisphosphate, and a fructose-2,6-biphosphatase activity that catalyzes the degradation of fructose-2,6-bisphosphate. This protein regulates fructose-2,6-bisphosphate levels in the heart, while a related enzyme encoded by a different gene regulates fructose-2,6-bisphosphate levels in the liver and muscle. This enzyme functions as a homodimer. Two transcript variants encoding two different isoforms have been found for this gene.

#### **Recommended Dilutions**

WB 1:1000 - 1:5000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID**5208

Swiss Prot
060825

#### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

#### **Synonyms**

PFK-2/FBPase-2; Phospho-PFKFB2-S483

#### Contact

www.abclonal.com


#### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

### **Validation Data**



Western blot analysis of lysates from HeLa cells, using Phospho-PFKFB2-S483 Rabbit pAb (AP0784) at 1:2000 dilution. HeLa cells were treated with IGF-1 (50 ng/ml) at 37°C for 30 minutes after serum-starvation overnight.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5s.