# ABclonal www.abclonal.com

## Phospho-NF-κB2-S866 Rabbit mAb

Catalog No.: AP1367 Recombinant

## **Basic Information**

## **Observed MW**

140kDa

#### **Calculated MW**

97kDa

#### Category

SMab Recombinant Monoclonal Antibody

## **Applications**

WB, ELISA

## **Cross-Reactivity**

Mouse

#### CloneNo number

ARC56344

## **Background**

This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NFkB). The NFkB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protein encoded by this gene can function as both a transcriptional activator or repressor depending on its dimerization partner. The p100 full-length protein is co-translationally processed into a p52 active form. Chromosomal rearrangements and translocations of this locus have been observed in B cell lymphomas, some of which may result in the formation of fusion proteins. There is a pseudogene for this gene on chromosome 18. Alternative splicing results in multiple transcript variants.

## **Recommended Dilutions**

WB

1:100 - 1:500

## **Immunogen Information**

Gene ID 4791 Swiss Prot Q00653

#### **Immunogen**

A synthetic phosphorylated peptide around S866 of human Phospho-NF- $\kappa$ B2-S866 (NP\_001070962.1).

#### **Synonyms**

p52; p100; H2TF1; LYT10; CVID10; LYT-10; NF-kB2; p49/p100; Phospho-NF-κB2-S866

## **Contact**

•

www.abclonal.com

## **Product Information**

**Source** Rabbit **Isotype** IgG Purification


Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of lysates from NIH/3T3 cells, using Phospho-NF- $\kappa$ B2-S866 Rabbit mAb (AP1367) at 1:500 dilution. NIH/3T3 cells were treated by TNF- $\alpha$  (20 ng/ml) at 37°C for 30 minutes.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.