

AP1399

Leader in Biomolecular Solutions for Life Science



Phospho-Jak1-Y1034/1035 Rabbit mAb

Catalog No.: AP1399

Recombinant

1 Publications

Basic Information

Observed MW

Refer to figures

Calculated MW

133kDa

Category

SMab Recombinant Monoclonal
Antibody

Applications

WB,ELISA

Cross-Reactivity

Human

CloneNo number

ARC3147

Recommended Dilutions

WB 1:500 - 1:1000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Contact

 www.abclonal.com

Background

This gene encodes a membrane protein that is a member of a class of protein-tyrosine kinases (PTK) characterized by the presence of a second phosphotransferase-related domain immediately N-terminal to the PTK domain. The encoded kinase phosphorylates STAT proteins (signal transducers and activators of transcription) and plays a key role in interferon-alpha/beta, interferon-gamma, and cytokine signal transduction. This gene plays a crucial role in effecting the expression of genes that mediate inflammation, epithelial remodeling, and metastatic cancer progression. This gene is a key component of the interleukin-6 (IL-6)/JAK1/STAT3 immune and inflammation response and is a therapeutic target for alleviating cytokine storms. The kinase activity of this gene is directly inhibited by the suppressor of cytokine signalling 1 (SOCS1) protein. Alternative splicing results in multiple transcript variants.

Immunogen Information

Gene ID

3716

Swiss Prot

P23458

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

JTK3; AIIDE; JAK1A; JAK1B; Phospho-Jak1-Y1022/1023

Product Information

Source

Rabbit

Isotype

IgG

Purification


Protein A purified

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PSB with 0.01% Sodium azide, 0.05% BSA, 40% glycerol, pH7.2.

Validation Data



Western blot analysis of lysates from Jurkat cells, using Phospho-Jak1-Y1034/1035 Rabbit mAb (AP1399) at 1:1000 dilution. Jurkat cells were treated with Pervanadate (1 mM) at 37°C for 30 minutes.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 60s.