

AP1462

Leader in Biomolecular Solutions for Life Science



Phospho-TIFA-T9 Rabbit mAb

Catalog No.: AP1462

Recombinant

Basic Information

Observed MW

21kDa

Calculated MW

21kDa

Category

SMab Recombinant Monoclonal
Antibody

Applications

WB, ELISA

Cross-Reactivity

Human

CloneNo number

ARC66052

Background

This gene encodes an adapter protein involved in adaptive and innate immunity. This protein includes a forkhead-associated (FHA) domain that specifically binds to phosphorylated serine and threonine residues. In response to bacterial infection, the encoded host cell protein undergoes an intermolecular interaction between the FHA domain and a phosphorylated threonine that leads to protein oligomerization and stimulation of the NF-kappa B and other downstream signaling pathways. This protein exhibits reduced expression in hepatocellular carcinoma and may suppress hepatocellular carcinoma progression. This protein may also play a role in the DNA damage response.

Recommended Dilutions

WB 1:2000 - 1:6000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

92610

Swiss Prot

Q96CG3

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

T2BP; T6BP; TIFAA; Phospho-TIFA-T9

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of lysates from TIFA knockout (KO) 293T cells transfected with TIFA or mutational TIFA, using Phospho-TIFA-T9 Rabbit mAb (AP1462) at 1:5000 dilution incubated overnight at 4°C. Human TIFA-transfected TIFA KO 293T cells were treated with ADP-heptose.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 45s.

WB samples for antibody validation are kindly provided by Dr. Feng Shao, NIBS