ABclonal www.abclonal.com

Recombinant Human IL-6 Protein

Catalog No.: RP00004 Recombinant 1 Publications

Sequence Information

Species Gene ID Swiss Prot <I>E. 3569 P05231 coli</I>

Tags

C-His

Synonyms

IL6;BSF-2;BSF2;CDF;HGF;HSF;IFNbeta-2;IFNB2;IL-6

Product Information

Source

<I>E. coli</I>

Purification

≥ 90 % as determined by SDS-PAGE;≥ 90 % as determined by HPI C.

Endotoxin

< 0.1 EU/µg of the protein by LAL method

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact



www.abclonal.com

Background

Interleukin-6 (IL-6) is a multifunctional α -helical cytokine that regulates cell growth and differentiation of various tissues, which is known particularly for its role in the immune response and acute phase reactions. The encoded protein has been shown to be an endogenous pyrogen capable of inducing fever in people with autoimmune diseases or infections. The protein is primarily produced at sites of acute and chronic inflammation, where it is secreted into the serum and induces a transcriptional inflammatory response through interleukin 6 receptor, alpha. The functioning of this protein is implicated in a wide variety of inflammation-associated disease states, including suspectibility to diabetes mellitus and systemic juvenile rheumatoid arthritis.

Basic Information

Description

Recombinant Human IL-6 Protein is produced by E. coli expression system. The target protein is expressed with sequence (Pro29-Met212) of human IL6 (Accession #NP_000591.1) fused with an initial Met at the N-terminus and a 6×His tag at the C-terminus.

Bio-Activity

1.Measured by its binding ability in a functional ELISA. Immobilized Recombinant Human IL6R at 1 μg/mL (100 μL/well) can bind Recombinant Human IL-6 with a linear range of 20-70 ng/mL.|2.Measured in a cell proliferation assay using TF-1 human erythroleukemic cells. The ED₅₀ for this effect is 0.18-0.74 ng/mL, corresponding to a specific activity of 1.35×10⁶~5.55×10⁶ units/mg..

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.


Avoid repeated freeze/thaw cycles.

^{*} For your safety and health, please wear a lab coat and disposable gloves when handling.


Validation Data




Recombinant Human IL-6 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



The purity of Human IL6 Protein (Cat.RP00004) was greater than 90% as determined by SEC-HPLC.



Immobilized Recombinant Human IL6R at 1 μ g/mL (100 μ L/well) can bind Recombinant Human IL-6 with a linear range of 20-70 ng/mL.



Recombinant Human IL6 stimulates cell proliferation of the TF-1 human erythroleukemic cells. The ED $_{50}$ for this effect is 0.18-0.74 ng/mL, corresponding to a specific activity of $1.35 \times 10^6 \sim 5.55 \times 10^6$ units/mg.