# ABclonal www.abclonal.com

# Recombinant Human IL-1 Beta/IL-1F2 Protein

Catalog No.: RP00454 Recombinant

# **Sequence Information**

Species Gene ID Swiss Prot <I>E. 3553 P01584 coli</I>

# Tags

C-His&Avi

#### **Synonyms**

Interleukin-1 beta; IL-1 beta; IL1F2; IL1B; IL-1BETA; IL1F2; IL-1 $\beta$ ; IL-1 beta; IL-1B ;Interleukin-1  $\beta$ ; IL-1  $\beta$ ; IL-1 $\beta$ ; IL-1 $\beta$ 

#### **Product Information**

**Source** Purification
HEK293 cells > 95% by SDS-

PAGE; > 95% by SEC-HPLC

SEC-H

#### **Endotoxin**

 $< 1 \; \text{EU/}\mu\text{g}$  of the protein by LAL method

#### **Formulation**

Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

#### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

# **Contact**

www.abclonal.com

# **Background**

Interleukin-1 beta (IL-1 $\beta$ ) is induced by inflammatory signals in a broad number of immune cell types. IL-1 $\beta$  (and IL-18) are the only cytokines which are processed by caspase-1 after inflammasome-mediated activation. IL-1 signaling activates innate immune cells including antigen presenting cells, and drives polarization of CD4 T cells towards T helper type (Th) 1 and Th17 cells.

### **Basic Information**

# **Description**


Recombinant Human IL-1 Beta/IL-1F2 Protein is produced by <I>E. coli</I> expression system. The target protein is expressed with sequence (Ala117-Ser269) of Human IL-1 Beta/IL-1F2 (Accession #P01584) fused with a C-His&Avi tag at the C-terminus.

#### **Bio-Activity**


#### Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. <br/> <br/> After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.


# **Validation Data**




Human IL-1 Beta on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Human IL-1 Beta is greater than 95% as determined by SEC-HPLC.



Immobilized Human IL-1 Beta, His Tag at 5  $\mu$ g/mL (100  $\mu$ L/Well) on the plate. Dose response curve for Human IL-1R2, hFc Tag with the EC<sub>50</sub> of 0.12  $\mu$ g/mL determined by ELISA (QC Test).



Human IL-1R2, hFc Tag captured on CM5 Chip via Protein A can bind Human IL-1 Beta, His Tag with an affinity constant of 0.88 nM as determined in SPR assay (Biacore T200).