

RP00465

Leader in Biomolecular Solutions for Life Science



Recombinant Human FGFR-3 beta (IIIc)/CD333 Protein

Catalog No.: RP00465 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
HEK293 cells	2261	P22607-1

Tags

C-His&Avi

Synonyms

ACH; CD333; CEK; CEK2; EC 2.7.10; FGF R3; FGFR3; HSGFR3EX; JTK4

Product Information

Source	Purification
Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.	≥ 95 % as determined by SDS-PAGE; ≥ 95 % as determined by HPLC.

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Background

Four distinct genes encoding closely related FGF receptors, FGF R1-4, are known. All four genes for FGF Rs encode proteins with an N-terminal signal peptide, three immunoglobulin (Ig)-like domains, an acid?box region containing a run of acidic residues between the IgI and IgII domains, a transmembrane domain and the split tyrosine-kinase domain. FGFR3 is tyrosine-protein kinase that acts as cell-surface receptor for fibroblast growth factors and plays an essential role in the regulation of cell proliferation, differentiation and apoptosis. Plays an essential role in the regulation of chondrocyte differentiation, proliferation and apoptosis, and is required for normal skeleton development. Regulates both osteogenesis and postnatal bone mineralization by osteoblasts.

Basic Information

Description

Recombinant Human FGFR-3 beta (IIIc)/CD333 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Asp127-Gly375) of Human FGFR3 beta (IIIc) (Accession #P22607-1) fused with a C-His&Avi tag at the C-terminus.

Bio-Activity

Shipping

Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.

Storage

Store at -20°C
Avoid repeated freeze/thaw cycles.

Reconstitution


Centrifuge the vial before opening.
Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact




www.abclonal.com


Validation Data



Recombinant Human FGFR-3 beta (IIIc)/CD333 Protein was determined by Tris-Bis PAGE under reducing conditions.



Immobilized Human FGFR3 beta (IIIc) at 0.5 µg/mL (100 µL/Well) on the plate. Dose response curve for Anti-FGFR3 Antibody, hFc Tag with the EC₅₀ of 8.5ng/mL determined by ELISA.



The purity of Human FGFR3 beta (IIIc) is greater than 95% as determined by SEC-HPLC.